CORRECTION Open Access



Correction: Impact of duration of treatments with metformin and sulfonylureas, individually or in combination, on diabetic retinopathy among newly diagnosed type 2 diabetic patients: a pooled cohort's analysis

Mansour Bahardoust¹, Yadollah Mehrabi¹, Farzad Hadaegh², Davood Khalili^{4*} and Ali Delpisheh^{1,3*}

Correction: Int J Retin Vitr 11, 9 (2025). https://doi.org/10.1186/s40942-025-00637-w.

Following publication of the original article [1], the authors reported the following errors:

The affiliation of Davood Khalili is incorrect. The correct affiliation is: Prevention of Metabolic Disorders Research Center, Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Further to this, it an error was reported in the Abstract, under the Methods heading. Tehran Sugar and Lipid Study (TLGS) should be Tehran Lipid and Glucose Study (TLGS).

The original article [1] has been corrected.

Published online: 12 March 2025

The online version of the original article can be found at https://doi.org/10.1186/s40942-025-00637-w.

*Correspondence:

Davood Khalili

dkhalili@endocrine.ac.ir

Ali Delpisheh

alidelpisheh@yahoo.com

¹Department of Epidemiology, School of Public Health & Safety, Shahid Beheshti University of Medical Sciences, Velenjak, 7th Floor, Bldg No.2 SBUMS, Arabi Ave, Tehran, Tehran Province, Iran

²Prevention of Metabolic Disorders Research Center, Division of Biostatistics and Epidemiology, Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, Velenjak, Yaman St, Aarabi St. No.24. Tehran. Iran

³Safety Promotion and Injury Prevention Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁴Prevention of Metabolic Disorders Research Center, Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran

References

 Bahardoust M, Mehrabi Y, Hadaegh F, et al. Impact of duration of treatments with Metformin and sulfonylureas, individually or in combination, on diabetic retinopathy among newly diagnosed type 2 diabetic patients: a pooled cohort's analysis. Int J Retin Vitr. 2025;11:9. https://doi.org/10.1186/s40942-02 5-00637-w.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/publiccommons.org/publiccommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.